



MAHAMUL03014/13/1/2012-TC

Volume: 11 Issue: 2 Month: June- Dec. 2023

ISSN:
Online : 2320-8341
Print : 2320- 6446

RESEARCH FRONT

An International Multidisciplinary Research Journal





MAHAMUL03014/13/1/2012-TC

Volume: 11 Issue: 2 Month: June- Dec 2023

RESEARCH FRONT

An International Multidisciplinary Research Journal

ISSN:
Online : 2320-8341
Print : 2320- 6446

Editorial Office

27B, Plot No. 17, Flat No. 5,
Shriramkunj Apartment
Yashwant Colony,
Camp Satara,
Maharashtra,
India.

Publisher

Mrs. M. S. Zodage
27B, Plot No. 17, Flat No. 5,
Shriramkunj Apartment
Yashwant Colony,
Camp Satara,
Maharashtra,
India.

Email

researchfront2013@gmail.com

Contact

Mobile:
09960544067
09561084067

Website

www.researchfront.org

Vol. 10, Issue - 3
(Sept. 2022 to Dec. 2022)

Chief Editor

Dr. S. B. Zodage

Dept. of Geography
Chhatrapati Shivaji College, Satara,
Maharashtra.

Executive Editors

Dr. L. Joe Morgan

Associate Professor
Jacksonville State University,
Jacksonville, Alabama, USA.

Dr. Bhim Prasad Kafle

Assistant Professor
Dept. of Natural Science,
Kathmandu University, Nepal

Dr. Narsimhulu Sanke

Assistant Professor
Dept. of Mechanical Engineering,
University College of Engineering,
Osmania University, Hyderabad.

Dr. S. T. Ingle

Professor & Director,
School of Environmental & Earth Sciences,
NM University, Jalgaon (MH)

Dr. D. B. Panaskar

Professor & Head,
Environmental Sciences,
SRTM University, Nanded (MH)

Dr. V. B. Zodage

Asso. Professor,
SRTM College, Kudal,
Dist - Sindhudurg (MH)

Editors

Dr. P. A. Khadke

Assistant Professor,
School of Earth Sciences,
SRTM University, Nanded, (MS)

Dr. V. R. Veer

Assistant Professor,
Kisan Veer Mahavidyalaya
Wai, Dist - Satara (MS)

Dr. S. M. Sathe

CSS College, Hupari,
Tal - Hatkangale, Dist - Kolhapur (MH)

Dr. S. V. Karande

Assistant Professor,
Chhatrapati Shivaji College, Satara (MH)

**CONTENTS**

1	Geographical Study of Origin, Migration of Banjara Community In Maharashtra State	1-7

The accountability of the research matter articulated in this journal is entirely of the author(s) concerned. The view expressed in the research papers/articles in this journal does not essentially correspond to the views of the publisher/editor. The publisher/editor of the journal is not liable for errors or any consequences arising from the exercise of information contained in it.

Geographical Study of Origin, Migration of Banjara Community In Maharashtra State

Dr. Rathod S. B.

Associate Professor & Head, Department of Geography,
Shri. R. R. Patil Mahavidyalaya, Savlaj Ta. Tasgaon Dist. Sangali

Abstract:

The study of crop combinations is a very significance aspect of agricultural geography. Crop combination defined that the total or aggregate of different crops cultivated/grown in a region at a given point of time. The crop combination technique represents the areas sharing significant proportion of crops at higher rank and it is a very necessary measurement to analyze of the cropping intensity. The physic-cultural conditions are responsible for variation in regional patterns of agricultural activity in an area. As a result there is marked regional disparities in land use pattern, cropping pattern, crop combination etc. in an areal unit at a point of time. Agriculture occupies an important status in Indian economy. The region having 10492 sq.km. area comprises 11 tahsils of Satara district. This study is based on secondary data collected from socio-economic abstract Satara district. An endeavour is made here to study the crop combination regions in Satara district for the year 2020-2021. The crop data has been computed with the help of Doi's technique of crop combination. Physiography, rainfall, soil and drainage influence impact on agricultural landuse pattern in district. Fourty one crops of have been considered for crop combination analysis. Among these cereals, cash crops, oil seeds, vegetable crops, fruit crops, fodder crops, flowers and Pulses are major crops. Crop combination, in Satara district has indentified four crop combination. Such type of study represents real situation of cropping pattern in Satara district and helps to planners and agricultural scientist for agricultural planning and tahsil level.

Keywords – Cropping Patterns, Crop Combination, Agricultural Development.

Introduction:

The study of crop combination regions constitutes important aspects of agricultural geography as it provides a good basic for agricultural regionalization. Crops are generally, grown in

combination in any region and the crops have its relative position in terms of crop combination. Farmers are growing numerous crops in the field rather than single crops. The distributional patterns of crops in any region are an outcome of predominance of certain or combination of crops. This is in term of emergence of typical crop combination. The statistical techniques provide accurate techniques for the study of agricultural landuse and cropping pattern various methods have used by scholars, scientists and planners. Weaver in 1954 has applied least standard deviation technique for crop combination regions. This method is based on the comparison of the actual percentage of cropped areas occupied by the different field crop with theoretical distribution is taken to formulate the crop combination. For a comprehensive and clear understanding of the agricultural mosaic of an agro-climatic region and for the planning and development of its agriculture, a systematic study of crop combinations is of great significance. The main advantage of the study of crop combinations regions lies in the fact that it enables to understand that impact of physical, and socio-economic conditions on the agricultural mosaic and scope for rotation and effect on double cropping, total production and per hectare productivity.


Study Area:

The Satara district is situated in west part in Maharashtra state. This district consists of eleven tahsils covering 1547 villages. The total area extent is of 10,492 square kilometer extending from 17°5' to 18°11' north latitude and 73°33' to 74°54' east longitudes. This district is confined by Pune district to north, Solapur district to east, Sangli district to south and Ratnagiri and Raigad district to west. Satara district has a typical landscapes due to variations in relief, climate and vegetation. The variation of relief ranges from the pinnacles and high plateau of the main Sahyadrians range having heights over 1500 meters above means sea level to the subdued basin of Nira river with an average height of about 600 meters above means sea level. The climate ranges from the rainiest in the Mahabaleshwar region which has an average annual of over 6000 mm to the driest in Man tahsil where the average annual rainfall is about 500 mm. The vegetal cover to varies from the typical monsoon forest in the west part to scrub and poor grass in the east parts. Jowar, Rice, Bajra, Sugercane, oil seeds and pulses are mainly cultivated in the district.

Objectives:

The main objectives of the present study are as follows,

1. To find out the crop combination regions in the study area.
2. To understanding the cropping pattern in different Tahsils of the district.



Research methodology:

The present study has been interpreted quantitatively and qualitatively with suitable research methods. Most of the data used in the paper is analyzed with the help of graphic representations, percentage representation and cartographic representation. The present study is based on secondary data collected from different published sources for the year 2020-2021. Tahsil wise land use data collected from socio-economic abstract, Satara district and district census handbook in Satara referred to collect related information. Topographical maps and survey of India sheets are used for physiographical study. Simple statistical method has used to compute the least sum

of squared deviation and variance and lowest standard deviation and coefficient of variation analysis (weaver) for cropping in the present study.

Crop combination analysis:

Crop combination is a dynamic concept. Cropping patterns and crop associations change in space and time. Recently the crop combination, analysis geographical studies has gained momentum and its important is increasing day by day. According to this method for present study in weavers crop combination (minimum deviation method) and calculating from 2018-2019 season data. The tahsil level crop combination has computed in this study. Table No. 2 shows combination like cereals, pulses, oilseeds, cash crops, fruits.

According to this method for present study in weaver crop combination (Minimum Deviation Method) and calculating from all seasons data (Table No. 2). The least sum of squared deviation and variance and lowest standard deviation and coefficient of variation formula was selected for this study and analysis from (2018-2019) seasons. Crops, vegetable crops, flower crops and fodder crops in the district. It is important to note that the environmental constraints owing to geographic situation, soil and climate have put the limit on diversified agricultural productivity. This is reflected in obtained resulted in the crop combination analysis.

Table No. 1: Crop combination region in Satara district

Sr. No.	Crop Combination	Name of Tahsil	Name of crop group
1	Monoculture	Khandala, Phaltan, Man, Khatav, Javali	Cereals
2	Two Crop Region	Koregaon, Satara Patan	Cereals + Pulses Cereals + oilseeds Cereals + oilseeds
3	Three Crop Region	Mahabaleshwar	Cereals + vegetable + fodder
4	Four crop region	Karad	Cereals + oilseeds + cash crops + fodder crops

Source: Computed by Authors (2022)

Devisions of Crop Combination:

1) Monoculture:

One crop group dominance or monoculture was practiced in Khandala, Phaltan, Man, Khatav and Javali Tahsil in Satara district. Jawar, bajara, rice, wheat and other cereals crops of the in tahsil. Jawar and bajara performs better in the arid and semi-arid areas. Rice grows well in the wet and warm areas in Javali tahsil. Jowar is cultivated in this region because in this region belonging to pore region. Rainfall is less in monsoon season.

2) Two crop combination region:

In this district two crop combination is found in Koregaon, Satara and Patan tahsil having combination of cereals, pulses and oilseeds crops groups. Cereals crops group is dominant crop are jawar, rice and wheat. Pulses crops group is dominant crop grown are Tur, and Oilseeds crops group is dominant crop grown are groundnut, sunflower, soyabean and kardai. Jawar, bajara and wheat are observed as dominant crop in Koregaon tahsil. The constituent crop of two crop combination, however, vary from region to region owing to great variations in the temperature and moisture conditions.

3) Three crop combination region :

The northern and western parts in the district in Mahabaleshwar tahsil have dominance of three crop group combination region. In this crops group combination are found cereals, vegetable and fodder crops group. Mahabaleshwar tahsil is situated in western part of Satara district. Such information both spatial and temporal is a prerequisite for understanding the agricultural potential of a region. Mahabaleshwar at top gets highest annual rainfall (6000 millimeters). Summers and quite warm while the winter are cool. Cereals, vegetable and fodder crops group is a combination which suits well to the prevailing agro-climatic conditions of the region.

4) Four crop combination region :

Four crops group combination is identified in Karad tahsil. Crops group like cereals, oilseeds, cash crop and fodder crops. The four crop combinational areas units of Karad tahsil grow jawar, groundnut, sugarcane, wheat and fodder crops in the Kharip and Rabi are the constituent crops. The cropping patterns of these region are likely to remain highly ramified unless provision of irrigation are developed or new seeds are developed which may suit to the prevailing environmental conditions of the region.




Figure No. 2: Satara District Crop Combination

Conclusion:

Weavers technique has identified four crops group combination in study region. Cereals as monoculture has found in Khandala, Phaltan, Man, Khatav and Javali tahsil. Jawar, bajara and wheat as monoculture has found in study region. Two crops group combination entered cereals and pulses, and cereals and oilseeds crop group. Three crops group combination appeared concentration in Mahabaleshwar tahsil. In this region cereals, vegetable and fodder crops group is most important and leading crop. Four crops group combination is identified in Karad tahsil. In this crop combination crops group like cereals, cash crop, oilseeds and fodder crops.

The agricultural development in Satara district is advancement, due to diversified nature of land use pattern and cropping pattern of the Satara district, has increased the production

capacity and cropping intensity of the land. In the present scenario need to strengthen the irrigation facilities, soil and moisture conservation, agronomic practices, changing in the cropping pattern, development of medium, adoption of bio-technology, afforestation, rural communications, livestock development, small and marginal farmers and agricultural laborers and setting up agro-based industries.

References:

1. Andrabi, Javaid Ahmad. Major crop combination and diversification regions in Kashmir Valley. Budapest International research in Exact Sciences Journal. 2019 January; 1(1): 54-61, ISSN: p-2655-7835.
2. Majid Husain (1999): “Systematic Agricultural Geography”, Rawat Public, New Delhi. Pp. 222 – 227.
3. Satara District Censes Handbook, 20011, District Socio-economic Abstract, Satara 2020.
4. Singh, R. P. (1967): Concept of land use, Patana University, Journal vol.22, pp. 52-62.
5. Das, M. M. (1984): Agricultural land use and cropping pattern in Assam, Geographical Review of India, Kolkata, vol.44/4.
6. Vaidya, B.C. (1990): Cropping pattern in Yashoda Basin, The Deccan Geographer, Secundarabad, 28/2 and 3.
7. Jasbir Sing, (1976): An Agricultural Geography of Haryana, Kurukshetra, Vishal Publishing Co.
8. Ayyar, N.P (1969): Crops Region of Madhya Pradesh: A Case Study in Methodology, Geography Review of India, Vol: 31, Pp.
9. Dhillan S. S. : Agricultural Geography.
10. Mamoria C. B. & Tripathi B. B. (1991), “Agricultural Problems of India”, Kitab Mahal, Allahabad.
11. Singh, R.P. (1967): Concept of landuse ,Patana University, Journal vol.22, pp.52-62.
12. Das, M.M.(1984): Agricultural landuse and cropping pattern in Assam, Geographical Review of India, Kolkata, vol.44/4.

MAHAMUL03014/13/1/2012-TC
Volume: 11 Issue: 2 Month: June - Dec. 2023

ISSN:
Online : 2320-8341
Print : 2320- 6446

